
Vanuatu

Total population (2016): 270 470 > Population aged 15 years and older (15+): 64% > Population in urban areas: 26% > Income group (World Bank): Lower-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	0.9		0.8	
Unrecorded	0.5		0.4	
Total**	1.1		1.0	
Total males / females	2.0	0.3	1.8	0.2
WHO Western Pacific Region	7.0		7	.3

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine

Spirits Other

	Litres
Males (15+)	8.9
Females (15+)	3.3
Both sexes (15+)	7.4

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	13.8	69.1	8.0	73.4
Females	2.5	33.7	1.4	37.2
Both sexes	8.1	59.5	4.8	64.8

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	54.6	80.4	67.6
Former drinkers* (15+)	25.5	12.2	18.8
Abstainers (15+), past 12 months	80.1	92.6	86.4

 $^{^{\}star}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF). $2016\,$

(1011), 2010					
	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	16.2	8.5	20.2	14.5	3
Road traffic injuries, males / females	26.0	12.0	16.1	10.5	5
Cancer, males / females	170.1	141.5	2.9	1.0	4

^{*}Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

^{*} Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	8.8	5.1
Females	1.8	0.7
Both sexes	5.2	2.9
WHO Western Pacific Region	4.7	2.3

 $^{^{\}star}$ 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2009/2015) / Yes
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	No / No / No
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	No / No / No
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, Yes / No, No No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	No legislation
Legally binding regulations on alcohol advertising / product placement (any)	No / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes